

Spontaneous emission of polaritons from a Bose-Einstein condensate

Karl-Peter Marzlin and Weiping Zhang

Department of Physics, Macquarie University, Sydney, NSW 2109, Australia

We study the spontaneous emission of a partially excited Bose-Einstein condensate composed of two-level atoms. The formation of polaritons induced by the ground-state part of the condensate leads to an avoided crossing in the photon spectrum. This avoided crossing acts similarly to a photonic band gap and modifies the spontaneous emission rate.

It is well-known that the radiation properties of atoms can dramatically be manipulated by changing the environment where the atoms emit photons. For microcavities it has been demonstrated [1] and for periodic dielectric media predicted [2,3] that the suppression of spontaneous emission (SE) can be quite high. In the case of a microcavity this happens because its geometry reduces the radiation-mode density, whereas in a periodic dielectric medium SE is suppressed due to the formation of photonic band gaps (PBG).

The recent achievement of Bose-Einstein condensation in magnetic traps [4] has provided a new state of matter where all atoms share a single macroscopic quantum state. Such a state of matter offers great opportunities to explore and test new phenomena related to macroscopic quantum coherence. Recently several authors have theoretically studied the optical properties of a Bose-Einstein condensate (BEC). For example, in a homogeneous, extended BEC polaritons (superpositions of excited atoms and photons) are formed [5]. In a spatially confined BEC, the continuous center-of-mass momentum distribution leads to an increase of SE [6,7]. In addition, the stimulated emission can be increased by the Bose enhancement in a BEC [8]. In the case of two BECs, interference effects can be important [9].

In the present paper we focus on the study on how the macroscopic quantum coherence of a ground-state homogeneous BEC can modify the radiation properties of atoms which are partially excited from the BEC by a coherent light field. In difference to a microcavity and PBG materials, the manipulation of the radiation properties in a homogeneously extended BEC is caused by the formation of polaritons due to the existence of ground-state macroscopic quantum coherence.

The model: We consider a BEC composed of two-level atoms which is coupled to the electromagnetic field. The coupling is described by using minimal coupling in rotating-wave approximation under neglect of the

term quadratic in the electromagnetic field. The interaction Hamiltonian is then given by

$$H_{\text{int}} = \int d^3k d^3k' \zeta_{\sigma}(\vec{k}) a_{\sigma}(\vec{k}) \Psi_g(\vec{k}') \Psi_e^{\dagger}(\vec{k} + \vec{k}') + \text{H.c.}, \quad (1)$$

with $\zeta_{\sigma}(\vec{k}) := \omega_0 \vec{d} \cdot \vec{\epsilon}_{\sigma}(\vec{k}) [\hbar / (2(2\pi)^3 \epsilon_0 \omega_k)]^{1/2}$ for a electromagnetic mode with polarization vector $\vec{\epsilon}_{\sigma}(\vec{k})$. The Heisenberg equations of motion for the photon annihilation operators $a_{\sigma}(\vec{k})$ and the field operators Ψ_e and Ψ_g for excited and ground-state atoms can be derived easily and are given by

$$i\hbar \dot{\Psi}_e(\vec{k}) = \left\{ \frac{\hbar^2 \vec{k}^2}{2M} + \hbar\omega_0 \right\} \Psi_e(\vec{k}) + \int d^3k' \sum_{\sigma} \zeta_{\sigma}(\vec{k}') a_{\sigma}(\vec{k}') \Psi_g(\vec{k} - \vec{k}') \quad (2)$$

$$i\hbar \dot{\Psi}_g(\vec{k}) = \frac{\hbar^2 \vec{k}^2}{2M} \Psi_g(\vec{k}) + \int d^3k' \sum_{\sigma} \zeta_{\sigma}^*(\vec{k}') a_{\sigma}^{\dagger}(\vec{k}') \Psi_e(\vec{k} + \vec{k}') \quad (3)$$

$$i\hbar \dot{a}_{\sigma}(\vec{k}) = \hbar\omega_k a_{\sigma}(\vec{k}) + \zeta_{\sigma}(\vec{k}) \int d^3k' \Psi_g^{\dagger}(\vec{k}') \Psi_e(\vec{k} + \vec{k}') \quad (4)$$

We have neglected the nonlinear interatomic interaction terms as they are small compared to the coupling energy $\hbar\sqrt{\omega_0 \nu_g}$, where we have defined $\nu_g := |\vec{d}|^2 \rho_g / (2\hbar\epsilon_0)$ with ρ_g being the density of atoms in the ground-state and $\vec{d}$ denoting the atomic dipole moment.

Macroscopic coherent solution: First, we consider the physics in the coherent regime where the atoms in the internal ground state share a single macroscopic wavefunction ψ_g and interact with a classical laser field $a_{\sigma}^{\text{class}}(\vec{k})$. This interaction prepares the BEC into a coherent superposition of the internal ground-state ψ_g and an excited state ψ_e . The equations of motion for these three collective fields are given by Eqs. (2) to (4) if the quantum field operators are replaced by the corresponding classical coherent fields.

We are interested in finding a particular solution of these equations corresponding to a BEC coherently coupled to a running laser wave. We thus make the ansatz $\psi_g(\vec{k}) = (2\pi)^{3/2} \sqrt{\rho_g} \delta(\vec{k}) \exp[-i\mu t]$ for a homogeneous ground-state BEC of density ρ_g , $a_{\sigma}^{\text{class}}(\vec{k}) = \exp[-i\omega_L t] \delta(\vec{k} - \vec{k}_L) \delta_{\sigma, \sigma_L} \Omega_L [2(2\pi)^3 \hbar \epsilon_0 \omega_{k_L}]^{1/2} / (|\vec{d}| \omega_0)$ corresponding to a laser beam with frequency ω_L , Rabi frequency $\Omega_L > 0$, polarization σ_L , and wavevector $\vec{k}_L$ (inside the

BEC), and $\psi_e(\vec{k}) = (2\pi)^{3/2} \sqrt{\rho_e} \delta(\vec{k} - \vec{k}_L) \exp[-i(\mu + \omega_L)t]$ describing coherent excited atoms of density ρ_e and of momentum $\hbar\vec{k}_L$. Inserting these expressions into the Heisenberg equations of motions leads to a set of algebraical conditions which fix the parameters μ , $\vec{k}_L$, and ρ_e which we assume to be smaller than ρ_g . If we neglect the kinetic energy the density of excited atoms is given by $\sqrt{\rho_e} = \sqrt{\rho_g} \Omega_L / (\mu + \Delta)$, where $\Delta := \omega_L - \omega_0$ is the detuning of the laser beam. The wavevector is fixed by $\omega_{k_L} = \omega_L/2 + \sqrt{(\omega_L/2)^2 - \omega_0^2 \nu_g / (\mu + \Delta)}$. For $\Delta \leq 0$ ($\Delta \geq 0$) the chemical potential is given by $\mu = -\Delta/2 \pm \sqrt{(\Delta/2)^2 + \Omega_L^2}$ which implies $\mu + \Delta > 0$ ($\mu + \Delta < 0$), respectively. Note that for $\Delta > 0$ the expression for ω_{k_L} implies the additional constraint $\mu + \Delta > 4\nu_g$.

Derivation of polariton modes: Having found the macroscopic coherent solution we now address the question of how quantum fluctuations will affect it. The corresponding stability analysis will determine how the SE process occurs as the system evolves away from the initial coherent solution. To perform this analysis we linearise Eqs. (2) to (4) around the coherent fields by inserting $\Psi_g = \exp[-i\mu t] \{\psi_g + \delta\Psi_g\}$ and $\Psi_e = \exp[-i(\mu + \omega)t] \{\psi_e + \delta\Psi_e\}$ as well as $a_\sigma = \exp[-i\omega t] \{a_\sigma^{\text{class}} + \delta a_\sigma\}$ and retaining only terms linear in $\delta\Psi_i$ and δa_σ , which describe the quantum fluctuations around the coherent solution.

The resulting linearized equations of motions can also be derived from an effective Hamiltonian for the quantum fluctuations,

$$H_{\text{fluct}} = H_{\text{pol}} + H_{\text{spont}}, \quad (5)$$

where

$$\begin{aligned} H_{\text{pol}} = & \hbar \int d^3k \left\{ \left[\frac{\hbar \vec{k}^2}{2M} - \Delta - \mu \right] \delta\Psi_e^\dagger \delta\Psi_e \right. \\ & + \left[\frac{\hbar \vec{k}^2}{2M} - \mu \right] \delta\Psi_g^\dagger \delta\Psi_g + \sum_\sigma (\omega_k - \omega_L) \delta a_\sigma^\dagger \delta a_\sigma \\ & + \Omega_L \left[\delta\Psi_g(\vec{k} - \vec{k}_L) \delta\Psi_e^\dagger(\vec{k}) + \text{H.c.} \right] \\ & \left. + \sum_\sigma \frac{\vec{d} \cdot \vec{\varepsilon}_\sigma}{|\vec{d}|} \sqrt{\frac{\nu_g \omega_0^2}{\omega_k}} \left[\delta a_\sigma(\vec{k}) \delta\Psi_e^\dagger(\vec{k}) + \text{H.c.} \right] \right\} \quad (6) \end{aligned}$$

describes the formation of polaritons and conserves the total number of particles. In contrast to previous work [5], it also includes an interaction with the ground-state fluctuations $\delta\Psi_g$ mediated through the laser field Ω_L . But since the interaction energy $\hbar \sqrt{\nu_g \omega_0^2 / \omega_k}$, (which we from now on will approximate by $\hbar \sqrt{\nu_g \omega_0}$) is much larger than $\hbar \Omega_0$ it will only give small corrections to the polariton energy. We therefore can neglect it. For the same reason we will omit the kinetic energy $\hbar^2 \vec{k}^2 / (2M)$ of the particles as well as the chemical potential μ .

The eigenmodes of H_{pol} are then easily found. To derive them it is of advantage to introduce the “non-

coupled” polarization vector $\vec{\varepsilon}_{\text{NC}}(\vec{p})$ which is perpendicular to the photon momentum $\vec{p}$ and the atomic dipole moment $\vec{d}$. Since the interaction couples the polarization to $\vec{d}$ a mode proportional to this vector describes uncoupled photons which do not contribute to SE. A second, “coupled”, polarization vector $\vec{\varepsilon}_C(\vec{p})$ is introduced by being perpendicular to $\vec{p}$ and $\vec{\varepsilon}_{\text{NC}}(\vec{p})$. It is easy to show that the scalar product $\vec{\varepsilon}_C(\vec{p}) \cdot \vec{d}$ is then given by $|\vec{d}| \sin \vartheta_{\vec{p}}$, where $\vartheta_{\vec{p}}$ is the angle between $\vec{p}$ and $\vec{d}$. We associate with $\vec{\varepsilon}_C(\vec{p})$ the quantum fluctuation operator $\delta a(\vec{k}) := \delta a_{\sigma=C}(\vec{k})$.

The relevant eigenmodes of H_{pol} then consist of two parts. (i) free ground-state atoms $\varphi_{g,\vec{k}}(\vec{p}) = \delta(\vec{p} - \vec{k})$ with momentum $\hbar\vec{k}$ and energy $E_{g,\vec{k}} \approx 0$, and (ii) polaritons with frequency spectrum

$$\omega_{\vec{k},\pm} = -\Delta \pm w_{\vec{k}}, \quad (7)$$

with $w_{\vec{k}} := \sqrt{(\Delta_k/2)^2 + \nu_g \omega_0 \sin^2 \vartheta_{\vec{k}}}$ and $\Delta_k := \omega_k - \omega_0$. The polariton modes are given by

$$\varphi_{\pm,\vec{k}}(\vec{p}) = \frac{\delta(\vec{p} - \vec{k})}{\sqrt{2w_{\vec{p}}}} \left(\pm \frac{\sqrt{\nu_g \omega_0} \sin \vartheta_{\vec{p}}}{\sqrt{w_{\vec{p}} \pm \Delta_{\vec{p}}/2}}, \quad \sqrt{w_{\vec{p}} \pm \Delta_{\vec{p}}/2} \right), \quad (8)$$

where the first component refers to $\delta\Psi_e(\vec{p})$ and the second one to $\delta a(\vec{p})$. The polariton spectrum $\omega_{\pm,\vec{p}}$ clearly exhibits an avoided crossing around $\Delta_{\vec{p}} = 0$ of width $\sqrt{\nu_g \omega_0} \sin \vartheta_{\vec{p}}$ (see Fig. 1). It also contains a small gap whose edge is reached in the limit $|\vec{p}| \rightarrow 0$ and ∞ [5].

The second part of the Hamiltonian,

$$H_{\text{spont}} = \hbar \omega_0 \sqrt{\nu_e} \int \frac{d^3k}{\sqrt{\omega_k}} \sin \vartheta_k \left\{ \delta a^\dagger(\vec{k}) \delta\Psi_g^\dagger(\vec{k}_L - \vec{k}) + \text{H.c.} \right\} \quad (9)$$

describes the spontaneous decay of excited atoms in the coherent field. It depends on the density of coherently excited atoms ρ_e through $\nu_e := \rho_e |\vec{d}|^2 / (2\hbar \varepsilon_0)$. Since it has a structure similar to the exponent of a squeezing operator we can expect that the coherent field configuration (corresponding to the vacuum plus an external field) is instable.

Spontaneous emission of polaritons: Having derived the polariton modes we are in the position to calculate the SE rate. This is done by expanding the field operators in terms of the eigenmodes,

$$(\delta\Psi_e(\vec{k}), \delta a(\vec{k})) = \int d^3p \sum_{n=\pm} \varphi_{n,\vec{p}}(\vec{k}) b_{n,\vec{p}}. \quad (10)$$

The operator $b_{n,\vec{p}}$ annihilates a polariton in the associated mode. In this base H_{pol} can be written as

$$H_{\text{pol}} = \int d^3p \sum_{n=\pm} \hbar \omega_{n,\vec{p}} b_{n,\vec{p}}^\dagger b_{n,\vec{p}} \quad (11)$$

The interaction hamiltonian H_{spont} which is responsible for SE can be written as

$$H_{\text{spont}} = \hbar\omega_0\sqrt{\nu_e} \int \frac{d^3k}{\sqrt{\omega_k}} \sum_{n=\pm} \sin\vartheta_k \sqrt{\frac{w_{\vec{k}} + n\Delta_{\vec{k}}/2}{2w_{\vec{k}}}} \times \left\{ b_{n,\vec{k}}^\dagger \delta\Psi_g^\dagger(\vec{k}_L - \vec{k}) + \text{H.c.} \right\}. \quad (12)$$

To derive the SE rate we have to determine the time evolution of polaritons around the coherent field configuration. We assume that initially all atoms and photons are described by the macroscopic coherent solution, or in other words, the quantized polariton field is initially in the vacuum $|0\rangle$. This state then evolves under the action of the polariton Hamiltonian. To describe this state $|\psi(t)\rangle$ we make the following ansatz, which corresponds to the one-photon approximation,

$$|\psi(t)\rangle \approx \alpha(t)|0\rangle + \int d^3k \sum_{n=\pm} \beta_{n,\vec{k}}(t) b_{n,\vec{k}}^\dagger \delta\Psi_g^\dagger(\vec{k}_L - \vec{k}) |0\rangle. \quad (13)$$

The Schrödinger equation $i\hbar|\dot{\psi}\rangle = H_{\text{fluct}}|\psi\rangle$ then can be solved by using the Laplace transform $\bar{\alpha}(s) = \int_0^\infty \exp[-ts]\alpha(t)dt$. In a standard manner [2] we then arrive at the equation for the Laplace transform,

$$\bar{\alpha}(s) = \alpha(0) [s + I(s)]^{-1}. \quad (14)$$

$I(s)$ can be written in the form

$$I = \frac{V\nu_e\omega_0^2}{(2\pi)^3} \int d^3k \frac{\sin^2\vartheta_{\vec{k}}}{\omega_{\vec{k}} \left(s - i\omega_L + i\omega_{\vec{k}} + \frac{\nu_g\omega_0 \sin^2\vartheta_{\vec{k}}}{s - i\Delta} \right)}, \quad (15)$$

where V denotes the quantization volume. This integral agrees with the one found in absence of a BEC (which describes SE in free space) by setting $\nu_g = \Delta = 0$. We denote this free space integral by $I_0 := I(\nu_g = \Delta = 0)$. Both integrals are linearly divergent and can be treated in the way pointed out by Bethe (see, e.g., Ref. [10]), i.e., we renormalize the integrals by subtracting the free-electron contribution,

$$I^{\text{Ren}} := I - \frac{V\nu_e\omega_0^2}{(2\pi)^3 i} \int d^3k \frac{\sin^2\vartheta_{\vec{k}}}{\omega_{\vec{k}}^2} \quad (16)$$

and $I_0^{\text{Ren}} = I^{\text{Ren}}(\nu_g = \Delta = 0)$. These renormalized integrals are only logarithmically divergent.

At this point it is customary in the calculation of the free-space SE to perform the Wigner-Weisskopf approximation by neglecting the dependence of $I_0^{\text{Ren}}(s)$ on s . In the presence of a band gap this is inappropriate due to the strong variation of the mode density around the gap [2,3]. Nevertheless, we can perform a generalized

Wigner-Weisskopf approximation in the following way. We expect that the typical timescale on which SE happens is much larger than the optical cycle timescale $1/\omega_0$. From the definition of the inverse Laplace transform,

$$\alpha(t) = \frac{1}{2\pi i} \int_{\epsilon - i\infty}^{\epsilon + i\infty} e^{ts} \bar{\alpha}(s) ds, \quad (17)$$

it is clear that the variable s plays more or less the role of a Fourier-transformed time. We thus expect that only values of s much smaller than ω_0 contribute significantly to the SE. This implies that we can neglect s wherever it appears together with ω_0 or ω_L . Thus, we are allowed to set $s - i\omega_L \approx -i\omega_L$ in the denominator of $I(s)$ while retaining the term depending on $s - i\Delta$.

In the case of I_0^{Ren} this procedure immediately reproduces the Wigner-Weisskopf result $I_0^{\text{Ren}} \approx N_e \{ (\gamma_0/2) + i\Delta_{\text{Lamb}}^{2\text{-lev}} \}$, where $\gamma_0 := \bar{d}^2\omega_0^3/(3\pi\hbar\epsilon_0 c^3)$ denotes the SE rate in free space and $N_e := V\rho_e$ the number of excited atoms in the BEC. To fix $\Delta_{\text{Lamb}}^{2\text{-lev}}$ we follow the theory of Bethe (see, e.g., [10]) and introduce a cut-off frequency of $m_e c^2/\hbar$ in I_0^{Ren} , where m_e is the electron's mass. Calculating the principal value of the integral then leads to $\Delta_{\text{Lamb}}^{2\text{-lev}} \approx 2\gamma_0$. In contrast to free space the SE rate in a BEC depends on $\Delta_{\text{Lamb}}^{2\text{-lev}}$ since such a radiative frequency correction shifts the center of the avoided crossing (or of a band gap [3]).

It remains to calculate a renormalized expression of the integral I^{Ren} in the presence of a BEC. Fortunately, this task reduces to integrals proportional to I_0^{Ren} and a couple of convergent integrals. Within the generalized Wigner-Weisskopf approximation we find

$$I^{\text{Ren}} = \left(1 + \frac{4i\nu_g}{5(s - i\Delta)} \right) I_0^{\text{Ren}} + \frac{N_e\gamma_0}{5\pi i} \left\{ \frac{47i\nu_g}{15(s - i\Delta)} + \frac{8}{3} - \frac{(s - i\Delta)}{i\nu_g} + \left(1 - \frac{4i\nu_g}{(s - i\Delta)} \right) \times \left(1 + \frac{(s - i\Delta)}{i\nu_g} \right)^{3/2} \text{arccoth} \left(\sqrt{1 + \frac{(s - i\Delta)}{i\nu_g}} \right) \right\} \quad (18)$$

To perform the inverse Laplace transformation we have to know the analytical structure of $\bar{\alpha}(s)$ of Eq. (14). Because of the complicated structure of Eq. (18) we eventually have to rely on numerical methods to determine $\alpha(t)$, but much can be gained by doing a general analysis first. In general, $\bar{\alpha}(s)$ has several poles and a branch cut originating from the term including the arccoth in I^{Ren} . This cut lies between $s = i\Delta$ and $s = i(\Delta - \nu_g)$. The inverse Laplace transform $\alpha(t)$ is then given by the residues of $\bar{\alpha}(s)$ at the poles s_i plus a contribution from the branch cut,

$$\alpha(t) = a_0 \left\{ \sum_{\text{poles}} e^{ts_i} \text{Res}(\bar{\alpha}(s), s \rightarrow s_i) \right.$$

$$+ \lim_{\epsilon \rightarrow 0} \int_{i\Delta}^{i(\Delta - \nu_g)} ds e^{ts} [\bar{\alpha}(s + \epsilon) - \bar{\alpha}(s - \epsilon)] \Big\} , \quad (19)$$

where we have used that the poles seem always to be of first order and only the straight parts of the integration around the branch cut do contribute. We see that to each pole corresponds a fraction of excited atoms whose decay rate is given by the real part of the pole s_i . The branch cut corresponds to a fraction of excited atoms which decays in a non-exponential way.

The magnitude of the SE modification is determined by the ratio of ν_g to $|s - i\Delta|$ which in free space is of the order of $\gamma_0 N_e$. Since for contemporary BECs ν_g is at best in the order of γ_0 SE is notably changed only if there are very few excited atoms N_e , which implies $|\Delta| \gg \Omega_L$. As discussed above, the macroscopic coherent solution implies in this case for $\Delta > 0$ the additional constraint $\Delta > 4\nu_g$. In Fig. 2 the real part of the two dominating poles s_1, s_2 is shown as a function of Δ for the case $N_e = 1$ and $\nu_g = 2.5\gamma_0$ (corresponding to an atom density ρ_g of $5 \times 10^{14} \text{ cm}^{-3}$). For $\Delta > 0$ a third pole appears with a very small negative decay rate ($< 10^{-3}\gamma_0$). The occurrence of negative decay rates is consistent within the range of validity of the linearized equations for the quantum fluctuations and may indicate the formation of an atom-photon bound state [11]. Obviously the change in the SE rate can be quite large for small $|\Delta|$. According to Eq. (19) the fraction of atoms belonging to the poles can be easily calculated by determining the residue at the poles. It turns out that the poles whose real part asymptotically approaches γ_0 always dominate and that the fraction of atoms belonging to other poles is significant only for small $|\Delta|$. The same holds for the fraction corresponding to the branch cut.

If $|\Delta| \gg \gamma_0$ holds the dominant pole s_1 can be calculated by perturbation theory. Its real part (the decay rate) is then given by

$$\frac{1}{2}\gamma(\Delta) = -N_e \frac{\gamma_0}{2} \left(1 - \frac{4\nu_g}{5\Delta} \right) . \quad (20)$$

We see that the SE rate is altered by a factor of $1 - 4\nu_g/(5\Delta)$. It depends on the sign of Δ whether SE is increased or decreased.

In conclusion we have shown that for $N_e = O(1)$ the spontaneous emission is suppressed by the presence of a BEC in the internal ground state. The latter induces an avoided crossing in the polariton spectrum which acts similarly to an effective band gap, thereby suppressing spontaneous emission.

Acknowledgement: This work has been supported by the Australian Research Council.

- [1] P. Goy, J.M. Raimond, M. Gross, and S. Haroche, Phys. Rev. Lett. **50**, 1903 (1983).
- [2] S. John and T. Quang, Phys. Rev. A **50**, 1764 (1994).
- [3] A.G. Kofman, G. Kurizki, and B. Sherman, J. Mod. Opt. **41**, 353 (1994).
- [4] M. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wieman, and E.A. Cornell, Science **269**, 198 (1995); C.C. Bradley, C.A. Sackett, J.J. Tollet and R. Hulet, Phys. Rev. Lett. **75**, 1687 (1995); M.-O. Mewes, M.R. Andrews, N.J. van Druten, D.M. Kurn, D.S. Durfee, C.G. Townsend and W. Ketterle, Phys. Rev. Lett. **77**, 416 (1996).
- [5] B.V. Svistunov and G.V. Shlyapnikov, Sov. Phys. JETP **71**, 71 (1990); H.D. Politzer, Phys. Rev. A **43**, 6444 (1991).
- [6] J. Javanainen, Phys. Rev. Lett. **72**, 2375 (1994).
- [7] L. You, M. Lewenstein, R.J. Glauber, and J. Cooper, Phys. Rev. A **53**, 329 (1996).
- [8] J.J. Hope and C.M. Savage, Phys. Rev. A **54**, 3177 (1996).
- [9] C.M. Savage, J. Ruostekoski, and D.F. Walls, Phys. Rev. A **56**, 2046 (1997).
- [10] P.W. Milonni, *The quantum vacuum*, Academic Press, San Diego 1994.
- [11] S. John and J. Wang, Phys. Rev. Lett. **64**, 2418 (1990).




FIG. 1. A BEC induces an avoided crossing in the polariton spectrum. Far away from the avoided crossing the polaritons describe excited atoms or photons. Thus, if one focuses on the photons, the avoided crossing provides an effective band gap.




FIG. 2. Spontaneous emission rate of a partially excited BEC for detuning $\Delta > 4\nu_g$ (solid lines) and $\Delta < 0$ (dashed lines). The quasi-particles break up into different fractions with different decay rates. The dominating fraction is the one whose decay rate asymptotically approaches γ_0 .